

TABLE 3. Treatment of Gonorrhea Infections Among Wisconsin Females Attending Family Planning, STD, and Dual-Protection Clinics, 2014

Jurisdiction of Client Provider Type	# Cases Reported	Cases Treated		Appropriate Regimen						
		#	%	Total Appropriate		Same Day		Within 14 Days		Within 30 Days
				#	%	#	%	#	%	#
Milwaukee County	368	317	86.1%	245	66.6%	65	17.7%	212	57.6%	236
Family Planning	254	212	83.5%	173	68.1%	27	10.6%	146	57.5%	166
STD	114	105	92.1%	72	63.2%	38	33.3%	66	57.9%	70
Dual Protection	0									
Other Wisconsin Counties	113	107	94.7%	84	74.3%	21	18.6%	71	62.8%	83
Family Planning	101	95	94.1%	72	71.3%	14	13.9%	60	59.4%	71
STD	8	8	100.0%	8	100.0%	4	50.0%	7	87.5%	8
Dual Protection	4	4	100.0%	4	100.0%	3	75.0%	4	100.0%	4
All Wisconsin Counties	481	424	88.1%	329	68.4%	86	17.9%	283	58.8%	319
Family Planning	355	307	86.5%	245	69.0%	41	11.5%	206	58.0%	237
STD	122	113	92.6%	80	65.6%	42	34.4%	73	59.8%	78
Dual Protection	4	4	100.0%	4	100.0%	3	75.0%	4	100.0%	4

30 Days

%

64.1%

65.4%

61.4%

73.5%

70.3%

100.0%

100.0%

66.3%

66.8%

63.9%

100.0%
