

TABLE 4. Treatment of Gonorrhea Infections Among Wisconsin Males Attending Family Planning, STD, and Dual-Protection Clinics, 2014

Jurisdiction of Client Provider Type	# Cases Reported	Cases Treated		Appropriate Regimen						
		#	%	Total Appropriate		Same Day		Within 14 Days		Within 30 Days
				#	%	#	%	#	%	#
Milwaukee County	597	561	94.0%	430	72.0%	334	55.9%	415	69.5%	424
Family Planning	202	185	91.6%	153	75.7%	113	55.9%	144	71.3%	150
STD	395	376	95.2%	277	70.1%	221	55.9%	271	68.6%	274
Dual Protection	0									
Other Wisconsin Counties	198	191	96.5%	159	80.3%	109	55.1%	150	75.8%	157
Family Planning	110	106	96.4%	79	71.8%	50	45.5%	71	64.5%	78
STD	42	39	92.9%	36	85.7%	26	61.9%	35	83.3%	35
Dual Protection	46	46	100.0%	44	95.7%	33	71.7%	44	95.7%	44
All Wisconsin Counties	795	752	94.6%	589	74.1%	443	55.7%	565	71.1%	581
Family Planning	312	291	93.3%	232	74.4%	163	52.2%	215	68.9%	228
STD	437	415	95.0%	313	71.6%	247	56.5%	306	70.0%	309
Dual Protection	46	46	100.0%	44	95.7%	33	71.7%	44	95.7%	44

30 Days

%

71.0%

74.3%

69.4%

79.3%

70.9%

83.3%

95.7%

73.1%

73.1%

70.7%

95.7%
